

Plastic responses to novel environments are biased towards phenotype dimensions with high additive genetic variation

Daniel W. A. Noble^{a,b,1,2}, Reinder Radersma^{c,1,2,3}, and Tobias Uller^c

^aDivision of Ecology and Evolution, Research School of Biology, The Australian National University, Canberra, ACT 2600, Australia; ^bEcology and Evolution Research Centre, School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW 2052, Australia; and ^cDepartment of Biology, Lund University, 223 62 Lund, Sweden

Edited by Marcus W. Feldman, Stanford University, Stanford, CA, and approved May 22, 2019 (received for review December 13, 2018)

Environmentally induced phenotypes have been proposed to initiate and bias adaptive evolutionary change toward particular directions. The potential for this to happen depends in part on how well plastic responses are aligned with the additive genetic variance and covariance in traits. Using meta-analysis, we demonstrate that plastic responses to novel environments tend to occur along phenotype dimensions that harbor substantial amounts of additive genetic variation. This suggests that selection for or against environmentally induced phenotypes typically will be effective. One interpretation of the alignment between the direction of plasticity and the main axis of additive genetic variation is that developmental systems tend to respond to environmental novelty as they do to genetic mutation. This makes it challenging to distinguish if the direction of evolution is biased by plasticity or genetic “constraint.” Our results therefore highlight a need for new theoretical and empirical approaches to address the role of plasticity in evolution.

phenotypic plasticity | phenotypic accommodation | cryptic genetic variation | evolvability

Populations that encounter novel environments must often adapt to persist. The rate and direction of adaptive change depends on the phenotypic and genetic variation that is exposed to natural selection. The “plasticity-first” hypothesis proposes that environmentally induced phenotypic variation initiates and directs adaptive genetic divergence along particular trajectories (1, 2). This has been demonstrated experimentally (3, 4), and the observation that plastic responses sometimes are aligned with population divergence in morphology, physiology, and behavior makes it a plausible explanation for adaptive divergence in the wild (5–10). However, whether or not evolutionary divergence generally is disposed to follow the direction of environmentally induced phenotypes is contested and poorly understood (reviewed in refs. 2 and 11–14).

Adaptation following exposure to novel or stressful environments is hypothesized to involve the refinement, exaggeration, or elimination of ancestrally plastic responses through quantitative genetic change. This is possible only when there is heritable variation along the dimensions of the phenotype that are plastic. In quantitative genetics, this variation is described in terms of additive genetic variance and covariance between traits, commonly referred to as the **G** matrix. The predicted change in phenotype under directional selection on standing genetic variation has been considered a quantitative genetic estimate of evolvability (15). Quantitative genetic evolvability is highest in the direction of phenotypic space with the most additive genetic variance. All else being equal, quantitative genetic evolvability of environmentally induced phenotypes will therefore be high when the direction of phenotypic plasticity is well aligned with the axis of maximum additive genetic variation in novel environments. We refer to this alignment as the short-term evolutionary potential (ref. 1, p. 142) of environmentally induced phenotypes or plastic responses.

There are reasons to believe that the direction of plasticity and the main axis of genetic variation can be aligned. Both genetic and environmental perturbation produce their phenotypic effects through development, and phenotypes that are induced by mutation can often also be induced by environmental factors, and vice versa (1). The **G** matrix is partly a result of the distribution of mutational effects (in addition to, e.g., selective loss of variation) and will therefore tend to reflect how traits are coregulated in development (16). Indeed, models demonstrate that correlational selection can produce developmental interactions that channel the phenotypic effects of mutation along dimensions of trait combinations that were favored in the past (17–21). Since phenotypic accommodation involves the coordinated adjustment of traits that need to function together (1), it too relies on developmental interactions that likely are products of correlational selection. Selecting for environmentally induced phenotypes may therefore change developmental regulation of phenotypes such that it increases the additive genetic variation along the dimensions of the phenotype that are plastic (22). If so, it implies that selection in the direction or opposition of plastic responses to novel environments will typically be effective, which may predispose the population to evolve in the short term along the phenotype dimensions that are plastic.

Significance

Theory suggests that populations can evolve by genetic modification of environmentally induced responses. We demonstrate that such plastic responses to novel environments tend to occur along trait dimensions that have high genetic variation. This suggests that selection for or against environmentally induced phenotypes typically will be effective. That organisms respond to environmental novelty and genetic variation in similar ways can be explained by how development is wired. This makes it challenging to distinguish if plasticity or genetic constraints shape evolutionary responses following environmental change.

Author contributions: D.W.A.N., R.R., and T.U. designed research; D.W.A.N., R.R., and T.U. performed research; D.W.A.N. and R.R. analyzed data; and D.W.A.N., R.R., and T.U. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Published under the PNAS license.

Data deposition: All raw matrix data, code, and analyses have been deposited in the Open Science Framework repository at <https://osf.io/fz7sr/>.

¹D.W.A.N. and R.R. contributed equally to this study.

²To whom correspondence may be addressed. Email: daniel.noble@anu.edu.au or reinder.radersma@wur.nl.

³Present address: Biometris, Wageningen University & Research, 6708 PB Wageningen, The Netherlands.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1821066116/-DCSupplemental.

Published online June 19, 2019.

Here we use a meta-analysis to test if plastic responses are biased toward trait dimensions that harbor high levels of additive genetic variation. Plastic responses can be described by a “plasticity vector” that quantifies the change in mean phenotypic trait values between two environments, expressed as the distance between the multivariate phenotype means. We chose to focus on environmental conditions to which the population is not (yet) adapted, since this is the most contentious scenario for plasticity-driven evolution. The potential for adaptive divergence along the trait dimensions that are environmentally responsive will be high when (i) genetic variation underlying phenotypic variance and covariance (the **P** matrix) increases in the novel environment and (ii) the plastic response (“plasticity vector”) falls along the main axis of **G**, while the orientation of **G** stays the same in the novel environment, or (iii) the plastic response does not fall along the main axis of **G**, but the orientation of **G** rotates in the novel environment such that it aligns with the direction of plasticity (Fig. 1). Our meta-analysis tests how common these scenarios are.

Results

We conducted a systematic search of the literature (see *Materials and Methods* for more detail) and identified 21 studies across 19 species where **G** and **P** matrices could be compared (i.e., 32 environmental comparisons of **G** and **P** with $n = 64$ effects total). We also identified 32 studies ($n = 98$ effects) across 30 species that could be used to derive estimates of the quantitative genetic evolvability in the direction of plasticity. This larger number of studies could be included because we only needed a vector of trait means and **G** matrices to generate effect sizes (see *Materials and Methods* for more detail). Data were taxonomically biased toward insects with a total of 56 effect sizes, whereas fish and plants had 18 and 16 effect sizes, respectively. There were also **G** matrices estimated for one snail and one frog species (having six and two effects, respectively). Matrix size

varied from 2 to 11 traits (mean = 4.16, $n = 98$), and the novel environments were imposed under conditions that were either “stressful” (resulting in decreased survival) or “nonstressful,” and using different breeding designs. Given these factors can affect **G** (23–25) we controlled for them in our analysis (see below and *Materials and Methods* for more details).

Changes in Additive Genetic Variance in Response to Novel Environments.

We first explored if novel environmental conditions result in an increase in additive genetic variation (i.e., “cryptic genetic variation”; refs. 23 and 24) by generating effect sizes that contrasted the structure of **G** between nonnovel and novel environments. Non-novel environments were those that the populations were considered to be well adapted to, or in which populations had been reared for at least five generations or more, whereas novel environments were those that were manipulated outside normal rearing conditions. We quantified changes in additive genetic variation by calculating a number of effect sizes describing changes in **G** across environments (see *Materials and Methods* for more detail). Here we focus on those describing the change in total standardized genetic variation (SDV) and the change in the proportion of additive genetic variation along the major axis of **G** (i.e., g_{max}) (PV_{max}) as these capture how heritable variation may be released in response to novel conditions. For some studies, novel environments were considered stressful and/or led to a decrease in survival (see *Materials and Methods*). Using a metaregression model we therefore tested whether the observed changes in **G** depended on whether the novel environment was “stressful” or “nonstressful.” We also tested whether the type of breeding design (“full-sib” vs. “half-sib”) and the number of traits impacted the magnitude of change.

Across studies, there was little evidence for a release of cryptic genetic variation (overall meta-analytic mean from intercept-only models: SDV = -0.12 , 95% CI = -0.47 to 0.23 ; PV_{max} = -0.06 , 95% CI = -0.68 to 0.55). However, there was a high amount of




Fig. 1. Alternative scenarios describing differences in the short-term evolutionary potential of environmentally induced phenotypes. Ellipses depict multivariate phenotypic (**P**, light color) and genotypic (**G**, dark color) variation for two traits (Trait 1 and Trait 2). Arrows connecting the centroids of nonnovel (green) and novel (orange) environments are the plasticity vectors that describe the change in multivariate mean phenotype (**P**) between two environments. (A) Plastic responses to novel environments can be said to have high evolutionary potential if there is ample genetic variation underlying **P** and the vector of plastic response is well-aligned with the major axis of genetic variation in the novel environment (i.e., dark orange ellipse). (B) Plastic phenotypic responses can be said to have low evolutionary potential if there is limited genetic variation underlying **P** and the vector of plastic response does not align with the major axis of genetic variation in the novel environment.




Fig. 3. Mean effect sizes comparing plastic responses to **G** in nonnovel and novel environments (Nonnovel vs. Novel) and using different breeding designs (Half-sib vs. Full-sib). (A) Proportion of genetic variation along the plastic response vector (π_e). (B) The alignment between the plastic response vector and main axes of genetic variation \mathbf{g}_{max} (θ_e). Negative values indicate that the plastic response vector and \mathbf{g}_{max} is fully aligned (i.e., $\sim 0^\circ$), whereas positive values indicate that the plastic response vector and \mathbf{g}_{max} is not aligned (i.e., $\sim 90^\circ$). Values of 0 indicate that the plastic response vector and \mathbf{g}_{max} are 45° to one another. Note that in all cases effect sizes are means for a study for an average number of traits and are independent of other factors (i.e., marginalized means) in the model (*Materials and Methods*).

It is not fully understood why genetic and environmental perturbations should have similar phenotypic effects. However, theoretical models have demonstrated that evolved regulatory systems tend to produce developmental biases that reflect correlational selection in the past (reviewed in refs. 20 and 26). For example, multivariate selection can predispose developmental systems to respond plastically in the direction of trait covariance (22). Similar biases in phenotypes can occur in response to mutation (e.g., ref. 21). The matrix of mutational variance and covariance (the **M** matrix; refs. 27 and 28) is a more appropriate estimate of developmental bias than **G** since the latter also reflects the effects of selection and drift. Nevertheless, **G** is often assumed to represent how phenotypes vary in response to genetic variation (e.g., refs. 29 and 30). In other words, since both the direction of plasticity and the **G** matrix capture how trait development is regulated, plastic responses and the major axis of additive genetic variation may commonly be aligned. Our results suggest that this may indeed be the case for many traits, even for plastic responses to environments to which organisms are not adapted.

Exposure to a novel or stressful environment may also be associated with a general increase in additive genetic variation (23). The release of “cryptic genetic variation” is sometimes considered central to plasticity-first models of evolution (2, 31). While additive genetic variation generally increased in novel environments, studies using half-sib breeding designs actually supported a decrease in additive genetic variance. This difference between breeding designs suggests that maternal or paternal effects contribute significantly to the phenotypic variation that is expressed in novel environments, and that the variance contributed by parental effects may result in better alignment with the plastic response. This result provides a more nuanced understanding of previous work; without effectively disentangling environmental and genetic effects studies may overestimate how much genetic variation is “cryptic” (32, 33).

While these results suggest substantial scope for population divergence in the direction of plasticity, the impact of plasticity on evolutionary trajectories depends on the strength and form of

selection operating on the phenotype distribution in novel environments—information that is often missing. For example, the studies in our meta-analysis rarely estimated selection on phenotypes in the novel environment (this also means we were unable to account for any selection that may have occurred due to mortality). This is not surprising given that estimating selection demands additional or more complex experiments. To date, only one study appears to have explicitly addressed the alignment between \mathbf{g}_{max} , plasticity, and selection (34). Lind et al. (34) showed that, in water fleas, adaptive plasticity in response to fish predator cues was well-aligned with \mathbf{g}_{max} and the response to survival selection; however, the same was not true in response to midge predatory cues. It is unclear if this difference is related to the strength of past selection, but the water flea clones were considered to be well-adapted to both predator regimes (which is why the study was not included in our meta-analysis).

That evolutionary responses are not easily predicted on the basis of either selection or variability is also evident by a compilation of 16 long-term study populations, which showed that, as environments change, the relationship between heritable variation in single traits (i.e., h^2) and selection is weak (35). Indeed, some of the studies in the present analysis found that the direction of plasticity and standing genetic variation were poorly aligned. On the basis of the theoretical models described above, it may appear reasonable to conclude that such instances of poor alignment simply reflect that individuals fail to successfully accommodate to novel conditions. However, this interpretation is complicated not the least by the fact that **G** also reflects selective removal of genetic variation (36). This means that most additive genetic variance detected by the studies in this meta-analysis could be selectively neutral in the ancestral environment. If so, the alignment between additive genetic variance and the direction of plasticity can be interpreted as if plasticity is biased toward dimensions of the phenotype that have been weakly, rather than strongly, selected in the past. This ambiguity in the interpretation of an alignment between plastic responses and **G** calls for




Fig. 4. (A) The total amount of genetic variation along the plastic response vector as a proportion of the maximum amount of variation in any direction, π_e . Smaller values on the axes (i.e., negative numbers) correspond to situations where genetic variation along the plasticity vector is low ("Low" label) whereas larger positive values correspond to situations where genetic variation along the plasticity vector is high ("High" label) for the nonnovel (green text) and novel (orange text) environments. (B) The angle between the plastic response vector and \mathbf{g}_{\max} (θ_e) in novel and nonnovel environments across all studies. Labels correspond to values on the axes where the plasticity vector does not align with the major axis of genetic variation, \mathbf{g}_{\max} (Poorly Aligned) or where \mathbf{g}_{\max} aligns with the plasticity vector (Well Aligned) in both the nonnovel (green text) and novel (orange text) environments. Axes are on the transformed scale to facilitate comparisons with analyses. Large negative values (e.g., -8) indicate that the plastic response vector and \mathbf{g}_{\max} is fully aligned (i.e., $\sim 0^\circ$), whereas large positive values (e.g., 8) indicate that the plastic response vector and \mathbf{g}_{\max} is not aligned (i.e., $\sim 90^\circ$). Values of 0 indicate that the plastic response vector and \mathbf{g}_{\max} are 45° to one another. Mean effect sizes and 95% sampling intervals are depicted.

theoretical and empirical studies that can demonstrate how developmental evolution shapes plasticity and standing genetic variation, the latter through phenotypic effects of mutation and selective removal of phenotypes.

Our findings have important implications for testing the plasticity-first hypothesis. Such tests are often done by comparing ancestral-descendent populations (2, 5, 6, 19, 37–39). Typically, studies identify a set of traits that have diverged between the populations and establish whether ancestral populations show plasticity in the direction of descendent populations for those traits. However, while the alignment between plastic responses and the major axis of genetic variation make plasticity-led divergence more likely, it also makes it challenging to distinguish signatures of plasticity-led evolution from evolutionary divergence biased by the phenotypic effects of genetic mutation (i.e., "genetic constraints"; refs. 29 and 30). More generally, quantitative genetic estimates are likely to provide limited insights into more long-term evolution (40, 41). Distinguishing the distinct empirical traces of environmentally and genetically induced developmental biases in evolution will thus require identification of appropriate "null" models (38). Ancestral-descendant comparisons in natural populations combined with theoretical modeling and experimental evolution studies will likely be necessary to assess the role of plasticity in adaptive diversification.

Materials and Methods

Literature Searches and Data Collection. We did two literature searches for studies containing either (i) 'G matrix*' AND 'P matrix*' AND 'environ*' or (ii) 'genetic covariance' AND 'phenotyp* covariance' AND 'environ*' in the title, abstract or keywords to find studies that estimated \mathbf{G} and \mathbf{P} matrices in multiple environments. In both searches we used Web of Science and Scopus (search date October 9, 2017), given that different search engines can yield different results (42). We restricted our subject areas to biology, plant sci-

ences, evolutionary biology, ecology, developmental biology, and agriculture. In addition, we extracted \mathbf{P} and \mathbf{G} correlation/covariance matrices from primary studies collated by previous meta-analyses exploring the environmental sensitivity of \mathbf{G} (33, 43) and a meta-analysis compiling environmental effects on additive genetic variance (32).

To be included in our meta-analysis, the primary study had to (i) conduct a quantitative genetic breeding design in two or more environments, where one environment was considered representative for the population or species (i.e., nonnovel), and one or more environments were novel, (ii) measure two or more phenotypic traits, and (iii) present \mathbf{G} and \mathbf{P} correlation/covariance matrices for these traits or \mathbf{G} along with trait means in both environments. We defined a novel environment as one where the source population had not been experienced in the recent past (more than five generations) or where the authors had clearly stated the novelty of the environment within the paper. We further classified novel environments as stressful if (i) authors clearly stated that the environment was stressful and/or (ii) a decreased survival in the novel environment was evident. We acknowledge that novel environments not deemed stressful by authors or where mortality data were not provided may also be considered stressful in many cases, but the distinction is nevertheless potentially important since selective mortality can influence estimates of genetic and phenotypic variation. Relevant studies deemed to meet the above criteria based on title and abstracts were screened by two of the authors (D.W.A.N. and R.R.). In some studies, only correlation matrices were provided. These studies were included if we could obtain additive genetic variance estimates for the set of traits so that correlation matrices could be converted to covariance matrices (necessary for generating effect sizes, see below). In instances where only part of the necessary data were available (e.g., only \mathbf{G} and not \mathbf{P} , missing sample sizes, no variance estimates, etc.) we estimated \mathbf{G} and \mathbf{P} from raw data if they were deposited from the authors in repositories or contacted authors directly for the relevant information. If this information could not be obtained we excluded the study, or in some cases the trait, from our analysis. See *SI Appendix, Fig. S1* for a full PRISMA diagram. All raw matrix data extracted from studies (44–75), code, and analyses can be found at <https://osf.io/fz7sr/>.




Fig. 5. Examples of studies (with two trait matrices) with high (A and B) or low (C and D) alignment between the direction of plastic responses and standing genetic variation. Arrows are plastic response vectors; large light ellipses are **P** matrices and smaller dark ellipses are **G** matrices. Colors correspond to the nonnovel (green) and novel (orange) environments. (A) Kasule (44) used the cotton stainer bug (*Dysdercus fasciatus*) to estimate quantitative genetic variation in fecundity and adult size under stressful conditions using a half-sib breeding design. (B) Blanckenhorn and Heyland (45) used a full-sib breeding design to estimate quantitative genetic variation for hind limb size and development time in the yellow dung fly (*Scathophaga stercoraria*) under stressful conditions. (C) Via and Connor (46) studied the flour beetle (*Tribolium castaneum*) using a half-sib breeding design to estimate quantitative genetic variation for pupal weight and development time under nonstressful conditions. (D) Kause and Morin (47) studied the sawfly (*Priophorus pallipes*) and estimated quantitative genetic parameters for development time and body size using a full-sib breeding design under stressful environmental conditions.

Moderator (Predictor) Variables. We collected a number of moderator variables which we a priori expected to explain variation in effect sizes. These included the number of traits quantified (i.e., total size of the **G** matrix), stressfulness of environment, and the study design used to estimate **G** matrices (i.e., half-sib or full-sib) which distinguished narrow-sense and broad-sense estimates of additive genetic variation, respectively. For meta-analyses involving comparison of **G** across environments, we also categorized effect sizes as having come from studies where the novel environment was considered stressful (i.e., stressful vs. nonstressful) and included this in the relevant analyses. However, it is important to note that our main aim is not to isolate specific environmental predictors of variation in effect size but to assess the extent to which the direction of plasticity and the main axis of additive genetic variance is aligned for populations exposed to a novel environment.

Matrix-Based Effect Sizes and Sampling Variance. We were specifically interested in comparing **G** and **P** matrices within and across nonnovel and novel environments and determining how stressfulness of the novel environment, trait number, and study design impacted the magnitude of changes in matrices. In addition, we were also interested in the degree to which plastic responses (i.e., the plasticity vectors) align with genetic variation (**G**) in nonnovel and novel environments. These questions require effect size estimates that can be used to compare various aspects of matrices and their alignments (details on effect size measures used are discussed below).

Estimation of **G** and **P** occurs with varying levels of precision (76), and any comparisons both within and between environments should attempt to account for the effects of sampling variance to avoid potentially erroneous conclusions.

We took a flexible meta-analytic approach to effect size and sampling variance estimation with matrices that allowed us to make use of traditional multilevel meta-analytic models, which more effectively weight studies based on their precision. Given that no single matrix effect size captures all aspects of how matrices can change, we used a series of alternative effect size measures (Fig. 6 A–D and described in detail below) and generated their corresponding sampling variances for comparing **G** and **P** matrices within and across environments (Fig. 6E). While we compared both **G** and **P** to themselves across environments, we focus on comparisons of **G** only. This is appropriate since **G** and **P** typically were very similar (see analysis code and results at <https://osf.io/fz7sr/>). Before simulations, any nonpositive-definite matrices (i.e., nonpositive eigen values) were “bent” to make them positive-definite, and we excluded any matrix row and column entry where trait genetic or phenotypic variances were zero or negative (33). This ensured that all matrices used for simulations were positive definite.

To estimate sampling variability for matrix-based effect sizes, we used Monte Carlo simulations where each positive definite covariance matrix (Σ), along with its respective sample size (number of families or sires for **G** and number of individuals for **P**), was used to generate 5,000 simulated datasets of each trait matrix from a paper. Sets of traits had the same covariance structure as the

total amount of variation was first calculated for each matrix as the sum of all diagonal elements of Σ (i.e., the trace of the covariance matrix). We then calculated the standardized difference in total variance between the non-novel and novel \mathbf{G} matrices (SDV_G) as

$$SDV = 2 \frac{tr(\Sigma_{Nov}) - tr(\Sigma_{NN})}{tr(\Sigma_{Nov}) + tr(\Sigma_{NN})},$$

where the $tr(\Sigma_{NN})$ and $tr(\Sigma_{Nov})$ are the traces of the covariance matrices in the nonnovel environment and novel environment, respectively. We did not take the absolute difference in variance because we were interested in, and also had specific predictions for, how variance should change in response to novel environmental conditions. Positive standardized effect sizes indicate that total variance in multivariate phenotypes from novel environments were larger than total variance in nonnovel environments.

Comparing proportion of variation along major axes of G . We also estimated the proportion of variation in G ($PV_{G_{\max}}$) that was along the major axis of additive genetic variation (i.e., along g_{\max}) for each environment (Fig. 6A). We quantified the proportion of total variance along g_{\max} as

$$PV_{max} = \frac{\lambda_{max}}{tr(\Sigma)},$$

where λ_{max} is the first eigenvalue (i.e., the amount of variation along the first eigenvector) and the $tr(\Sigma)$ is the trace of the covariance matrix. Given that PV_{max} is a proportion, we logit-transformed it to normalize its distribution and sampling error.

Comparing major axes of genetic and phenotypic variation. Second, we explored how the orientations of **G** and **P** change within and across environments by calculating the angle between the major axes of variation (Fig. 6B). The major axis of variation for a covariance matrix is described by its first eigenvector. To compare within environments, we calculated the angle between \mathbf{g}_{\max} and \mathbf{p}_{\max} (θ_{NN} or θ_{Nov} depending on the environment), which are the first eigenvectors of **G** and **P**, respectively. To compare across the environments, we calculated the angle between the nonnovel \mathbf{g}_{\max} and novel \mathbf{g}_{\max} (θ_G), or the angle between the nonnovel \mathbf{p}_{\max} and novel \mathbf{p}_{\max} (θ_P), depending on whether the angle was calculated for the genetic or phenotypic variation. The angle between the two first eigenvectors was calculated as

$$\theta = \frac{180}{\pi} \cos^{-1} \left(\frac{V_{1_max} \cdot V_{2_max}}{\sqrt{V_{1_max} \cdot V_{1_max}} \cdot \sqrt{V_{2_max} \cdot V_{2_max}}} \right),$$

where v_{1_max} is the first eigenvector of one of the covariance matrices and v_{2_max} is the first eigenvector of the other covariance matrix. This calculation results in angles varying between 0° and 180° and so we subtracted from 180° any angle between 180° and 90° to yield angles between 0° and 90° (33). To normalize the distribution of angles we divided them by 90 and performed a logit transformation.

Assessing short-term evolutionary potential of plastic responses. Finally, to investigate the short-term evolutionary potential along the dimensions of plastic responses we tested whether or not the plastic response vector—the change in mean trait values between the nonnovel to novel environment—was aligned with the **G** matrices of the nonnovel and the novel environment. First, we calculated the vector $\Delta\bar{x}$ describing the change in mean trait values:

$$\Delta \bar{X} = \bar{X}_{NN} - \bar{X}_{Nov.}$$

Next, we projected the \mathbf{G} matrices for both the nonnovel and novel environments on the vector $\Delta\bar{\mathbf{x}}$ to calculate the amount of variance in \mathbf{G} in the direction of vector $\Delta\bar{\mathbf{x}}$ (i.e., $\mathbf{e}_{\Delta\bar{\mathbf{x}}}$) using equation 1 in ref. 15:

$$e_{\Delta\bar{x}} = \frac{(\Delta\bar{x})' G \Delta\bar{x}}{|\Delta\bar{x}|^2}.$$

To estimate how much variation was present in the direction of the plastic response compared with the maximum amount of variation in any direction, we divided $e_{\Delta\bar{x}}$ by the first eigenvalue of \mathbf{G} (λ_{G_max}), to calculate the evolutionary potential of plasticity (Fig. 6C):

$$\pi_e = \frac{e_{\Delta \bar{x}}}{\lambda_{G \max}}.$$

Given that π_e is a proportion, we logit-transformed it to normalize its distribution and sampling error. To test whether π_e was significantly larger than expected for any random alignment between the plastic response and

G, we also calculated a null expectation for the evolutionary potential of plasticity (π_0):

$$\pi_0 = \frac{\bar{\lambda}_G}{\lambda_{G, \max}},$$

in which $\bar{\lambda}_G$ is the average of all eigenvalues for \mathbf{G} . Next, we subtracted the logit of π_0 from the logit of π_e to estimate the difference between the observed and expected evolutionary potential in the direction of plasticity (π_{e-0}):

$$\pi_{e=0} = \text{logit}(\pi_e) - \text{logit}(\pi_0).$$

When the lower boundary of the credible interval for the intercept of the meta-analytic model (as described below) for $\pi_{e=0}$ is larger than zero, the evolutionary potential in the direction of plasticity is higher than what we would expect for a random alignment between $\Delta \bar{x}$ and \mathbf{g}_{\max} . We also calculated the angle between $\Delta \bar{x}$ and the first eigenvector of \mathbf{G} ($\mathbf{v}_{\mathbf{G}, \max}$) (Fig. 6D):

$$\theta_e = \frac{180}{\pi} \cos^{-1} \left(\frac{\Delta \bar{X} \cdot V_{G_max}}{\sqrt{\Delta \bar{X} \cdot \Delta \bar{X}} \cdot \sqrt{V_{G_max} \cdot V_{G_max}}} \right).$$

This calculation results in angles varying between 0° and 180° and so we subtracted from 180° any angle between 180° and 90° to yield angles between 0° and 90°. To normalize the distribution of angles we divided them by 90 and then performed a logit transformation.

Meta-Analysis. We meta-analyzed effect sizes using multilevel meta-analytic and meta-regression models (78) with the R package metafor (79). In all models, effects were weighted by their sampling variance from the simulations and we included “study” as a random effect. For analyses comparing environments, each study contributed multiple estimates and so we included a random slope of environment type in the study-level random effect to account for this nonindependence. However, only a random intercept was included if variance in the slope failed to converge in the model. Given that there was strong overlap between study and species we did not include a species-level random effect or phylogeny in our models to avoid confounding estimates (80). The package metafor does not estimate a residual variance by default so we included an observation-level random effect in all our models. Multiple environments were manipulated in some studies, and given quantitative genetic designs often separated families across these environments, this introduces a level of nonindependence between effect sizes (80). Given the limited sample sizes, we did not account for this level of dependence; however, we acknowledge this may lead to inflated type I error rates.

We used intercept only meta-analytic models to quantify total (I_t^2) and between study (I_{st}^2) heterogeneity in effects (81, 82) as

$$I_{st}^2 = \frac{\sigma_{st}^2}{\sigma_{st}^2 + \sigma_a^2 + \sigma_m^2}$$

$$I_t^2 = \frac{\sigma_{st}^2 + \sigma_e^2}{\sigma_{st}^2 + \sigma_e^2 + \sigma_m^2}$$

$$\sigma_m^2 = \frac{\sum w_i(k-1)}{[(\sum w_i)^2 - \sum w_i^2]},$$

where σ_m^2 is an estimate of the total sampling variability derived from w_i , the weight or inverse of the sampling variance, and k the total number of effects. σ_{τ}^2 and σ_e^2 are the between- and within-study variance estimates. I_{τ}^2 can be interpreted as the proportion of variation in effects resulting from differences between studies, which includes differences such as species, trait types, and methodological approaches that vary across studies. In contrast, I_{τ}^2 is the total proportion of variation in effects beyond sampling variance. If I_{τ}^2 is high, then this suggests that the effects observed are not simply the result of sampling variability.

We ran metaregression models to explore how our predicted moderator variables explained variation in effects. For each effect size, we included main effects for moderators we hypothesized would explain variation in effects when sample sizes allowed. These included (i) the number of traits, (ii) environment type (nonnovel and novel—only for analyses comparing the alignment between the plasticity vector and **G**), (iii) the type of novel environment (i.e., stressful or nonstressful—only for analyses comparing **G** across environments), and (iv) the breeding design (i.e., half-sib or full sib). Given the limited sample sizes and uneven taxonomic sampling we restricted models to estimating main effects of the above moderators only. Models comparing **G**

and P across environments were run separately. We calculated the marginalized mean effect size (and confidence intervals) for different subgroups of categorical predictors by taking the weighted average of the mean (and sampling variance) of one level in factor 1 (e.g., half-sib for the breeding design factor) across both levels of factor 2 (i.e., stressed and nonstress for the novel environment factor). We did this for all combinations to produce mean estimates unconditional on other categorical factors in the model. We checked funnel plot (i.e., effect size as a function of sampling error) asymmetry to look for evidence of publication bias. However, we found no clear evidence for publication bias in any of the effect sizes (*SI Appendix, Fig. S2*).

1. M. J. West-Eberhard, *Developmental Plasticity and Evolution* (Oxford University Press, New York, 2003).
2. N. A. Levis, D. W. Pfennig, Evaluating 'plasticity-first' evolution in nature: Key criteria and empirical approaches. *Trends Ecol. Evol.* **31**, 563–574 (2016).
3. Y. Suzuki, H. F. Nijhout, Evolution of a polyphenism by genetic accommodation. *Science* **311**, 650–652 (2006).
4. C. H. Waddington, Canalization of development and genetic assimilation of acquired characters. *Nature* **183**, 1654–1655 (1959).
5. N. A. Levis, A. J. Isdane, D. W. Pfennig, Morphological novelty emerges from pre-existing phenotypic plasticity. *Nat. Ecol. Evol.* **2**, 1289–1297 (2018).
6. A. G. Scoville, M. E. Pfrender, Phenotypic plasticity facilitates recurrent rapid adaptation to introduced predators. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 4260–4263 (2010).
7. I. Gomez-Mestre, D. R. Buchholz, Developmental plasticity mirrors differences among taxa in spadefoot toads linking plasticity and diversity. *Proc. Natl. Acad. Sci. U.S.A.* **103**, 19021–19026 (2006).
8. M. A. Wund, J. A. Baker, B. Clancy, J. L. Golub, S. A. Foster, A test of the "flexible stem" model of evolution: Ancestral plasticity, genetic accommodation, and morphological divergence in the threespine stickleback radiation. *Am. Nat.* **172**, 449–462 (2008).
9. A. V. Badyaev, Evolutionary significance of phenotypic accommodation in novel environments: An empirical test of the Baldwin effect. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **364**, 1125–1141 (2009).
10. S. Casasa, A. P. Moczek, The role of ancestral phenotypic plasticity in evolutionary diversification: Population density effects in horned beetles. *Anim. Behav.* **137**, 53–61 (2018).
11. C. D. Schlichting, M. A. Wund, Phenotypic plasticity and epigenetic marking: An assessment of evidence for genetic accommodation. *Evolution* **68**, 656–672 (2014).
12. A. P. Moczek et al., The role of developmental plasticity in evolutionary innovation. *Proc. Biol. Sci.* **278**, 2705–2713 (2011).
13. C. K. Ghalambor, J. K. McKay, S. P. Carroll, D. N. Reznick, Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Funct. Ecol.* **21**, 394–407 (2007).
14. D. W. Pfennig et al., Phenotypic plasticity's impacts on diversification and speciation. *Trends Ecol. Evol.* **25**, 459–467 (2010).
15. T. F. Hansen, D. Houle, Measuring and comparing evolvability and constraint in multivariate characters. *J. Evol. Biol.* **21**, 1201–1219 (2008).
16. J. M. Cheverud, Quantitative genetics and developmental constraints on evolution by selection. *J. Theor. Biol.* **110**, 155–171 (1984).
17. K. Kouvaris, J. Clune, L. Kounios, M. Brede, R. A. Watson, How evolution learns to generalise: Using the principles of learning theory to understand the evolution of developmental organisation. *PLoS Comput. Biol.* **13**, e1005358 (2017).
18. M. Parter, N. Kashtan, U. Alon, Facilitated variation: How evolution learns from past environments to generalize to new environments. *PLoS Comput. Biol.* **4**, e1000206 (2008).
19. M. Pavlicev, E. A. Norgard, G. L. Fawcett, J. M. Cheverud, Evolution of pleiotropy: Epistatic interaction pattern supports a mechanistic model underlying variation in genotype-phenotype map. *J. Exp. Zool. B Mol. Dev. Evol.* **316**, 371–385 (2011).
20. T. Uller, A. P. Moczek, R. A. Watson, P. M. Brakefield, K. N. Laland, Developmental bias and evolution: A regulatory network perspective. *Genetics* **209**, 949–966 (2018).
21. R. A. Watson, G. P. Wagner, M. Pavlicev, D. M. Weinreich, R. Mills, The evolution of phenotypic correlations and "developmental memory". *Evolution* **68**, 1124–1138 (2014).
22. J. A. Draghi, M. C. Whitlock, Phenotypic plasticity facilitates mutational variance, genetic variance, and evolvability along the major axis of environmental variation. *Evolution* **66**, 2891–2902 (2012).
23. G. Gibson, I. Dworkin, Uncovering cryptic genetic variation. *Nat. Rev. Genet.* **5**, 681–690 (2004).
24. A. A. Hoffmann, J. Merilä, Heritable variation and evolution under favourable and unfavourable conditions. *Trends Ecol. Evol.* **14**, 96–101 (1999).
25. M. Lynch, B. Walsh, *Genetics and Analysis of Quantitative Traits* (Oxford University Press, Oxford, UK, 1998), p. 980.
26. R. A. Watson, E. Szathmáry, How can evolution learn? *Trends Ecol. Evol.* **31**, 147–157 (2016).
27. J. M. Cheverud, Phenotypic, genetic and environmental morphological integration in the cranium. *Evolution* **36**, 499–516 (1982).
28. A. G. Jones, S. J. Arnold, R. Bürger, The mutation matrix and the evolution of evolvability. *Evolution* **61**, 727–745 (2007).
29. D. Schluter, Adaptive radiation along genetic lines of least resistance. *Evolution* **50**, 1766–1774 (1996).
30. J. W. McGlothlin et al., Adaptive radiation along a deeply conserved genetic line of least resistance in Anolis lizards. *Evol. Lett.* **2**, 310–322 (2018).
31. R. Lande, Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. *J. Evol. Biol.* **22**, 1435–1446 (2009).
32. P. K. Rowinski, B. Rogell, Environmental stress correlates with increases in both genetic and residual variances: A meta-analysis of animal studies. *Evolution* **71**, 1339–1351 (2017).
33. C. W. Wood, E. D. Brodie, 3rd, Environmental effects on the structure of the G-matrix. *Evolution* **69**, 2927–2940 (2015).
34. M. I. Lind, K. Yarle, J. Reger, M. J. Carter, A. P. Beckerman, The alignment between phenotypic plasticity, the major axis of genetic variation and the response to selection. *Proc Biol Sci* **282**, 20151651 (2015).
35. J. J. C. Ramakers, A. Culina, M. E. Visser, P. Gienapp, Environmental coupling of heritability and selection is rare and of minor evolutionary significance in wild populations. *Nat. Ecol. Evol.* **2**, 1093–1103 (2018).
36. R. Lande, The genetic covariance between characters maintained by pleiotropic mutations. *Genetics* **94**, 203–215 (1980).
37. A. Corl et al., The genetic basis of adaptation following plastic changes in coloration in a novel environment. *Curr. Biol.* **28**, 2970–2977.e7 (2018).
38. K. Kovaka, Underdetermination and evidence in the developmental plasticity debate. *Br. J. Philosophical Biol.* **70**, 127–152 (2019).
39. R. J. S. McCairns, L. Bernatchez, Adaptive divergence between freshwater and marine sticklebacks: Insights into the role of phenotypic plasticity from an integrated analysis of candidate gene expression. *Evolution* **64**, 1029–1047 (2010).
40. T. F. Hansen, "Adaptive landscapes and macroevolutionary dynamics" in *The Adaptive Landscape in Evolutionary Biology*, E. Svensson, R. Calsbeek, Eds. (Oxford University Press, Oxford, UK, 2012), pp. 205–226.
41. A. G. Jones, S. J. Arnold, R. Bürger, Stability of the G-matrix in a population experiencing pleiotropic mutation, stabilizing selection, and genetic drift. *Evolution* **57**, 1747–1760 (2003).
42. P. Mongeon, A. Paul-Hus, The journal coverage of Web of Science and Scopus: A comparative analysis. *Scientometrics* **106**, 213–228 (2016).
43. C. W. Wood, E. D. Brodie, 3rd, Evolutionary response when selection and genetic variation covary across environments. *Ecol. Lett.* **19**, 1189–1200 (2016).
44. F. K. Kasule, Associations of fecundity with adult size in the cotton stainer bug *Dysdercus fasciatus*. *Heredity* **66**, 281–286 (1991).
45. W. U. Blanckenhorn, A. Heyland, The quantitative genetics of two life history trade-offs in the yellow dung fly in abundant and limited food environments. *Evol. Ecol.* **18**, 385–402 (2004).
46. S. Via, J. Conner, Evolution in heterogeneous environments: Genetic variability within and across different grains in *Tribolium castaneum*. *Heredity* **74**, 80–90 (1995).
47. A. Kause, J. P. Morin, Seasonality and genetic architecture of development time and body size of the birch feeding sawfly *Priophorus pallipes*. *Genet. Res.* **78**, 31–40 (2001).
48. J. R. Auld, The effects of predation risk on mating system expression in a freshwater snail. *Evolution* **64**, 3476–3494 (2010).
49. T. D. Beacham, A genetic analysis of early development in pink (*Oncorhynchus gorbuscha*) and chum salmon (*Oncorhynchus keta*) at three different temperatures. *Genome* **30**, 89–96 (1988).
50. M. Bégin, D. A. Roff, An analysis of G matrix variation in two closely related cricket species, *Gryllus firmus* and *G. pennsylvanicus*. *J. Evol. Biol.* **14**, 1–13 (2001).
51. M. Bégin, D. A. Roff, V. Debat, The effect of temperature and wing morphology on quantitative genetic variation in the cricket *Gryllus firmus*, with an appendix examining the statistical properties of the Jackknife-MANOVA method of matrix comparison. *J. Evol. Biol.* **17**, 1255–1267 (2004).
52. M. T. Brock et al., Floral genetic architecture: An examination of QTL architecture underlying floral (co)variation across environments. *Genetics* **186**, 1451–1465 (2010).
53. R. D. Collins, Y. Jang, K. Reinhold, M. D. Greenfield, Quantitative genetics of ultrasonic advertisement signalling in the lesser waxmoth *Achroia grisella* (Lepidoptera: Pyralidae). *Heredity* **83**, 644–651 (1999).
54. J. K. Conner, R. Franks, C. Stewart, Expression of additive genetic variances and covariances for wild radish floral traits: Comparison between field and greenhouse environments. *Evolution* **57**, 487–495 (2003).
55. M. E. Czesak, C. W. Fox, Evolutionary ecology of egg size and number in a seed beetle: Genetic trade-off differs between environments. *Evolution* **57**, 1121–1132 (2003).
56. M. Delcourt, M. W. Blows, H. D. Rundle, Sexually antagonistic genetic variance for fitness in an ancestral and a novel environment. *Proc. Biol. Sci.* **276**, 2009–2014 (2009).
57. M. Delcourt, M. W. Blows, H. D. Rundle, Quantitative genetics of female mate preferences in an ancestral and a novel environment. *Evolution* **64**, 2758–2766.
58. L. Engqvist, Environment-dependent genetic correlations between development time and body mass in a scorpionfly. *Zoology (Jena)* **110**, 344–353 (2007).

59. J. P. Evans, M. M. Rahman, C. Gasparini, Genotype-by-environment interactions underlie the expression of pre- and post-copulatory sexually selected traits in guppies. *J. Evol. Biol.* **28**, 959–972 (2015).
60. C. P. Grill, A. J. Moore, E. D. Brodie, III, The genetics of phenotypic plasticity in a colonizing population of the ladybird beetle, *Harmonia axyridis*. *Heredity* **78**, 261–269 (1997).
61. J. Guan *et al.*, Estimating genetic parameters and genotype-by-environment interactions in body traits of turbot in two different rearing environments. *Aquaculture* **450**, 321–327 (2016).
62. J. Guntrip, R. M. Sibly, G. J. Holloway, The effect of novel environment and sex on the additive genetic variation and covariation in and between emergence body weight and development period in the cowpea weevil, *Callosobruchus maculatus* (Coleoptera, Bruchidae). *Heredity* **78**, 158–165 (1997).
63. G. J. Holloway, S. R. Povey, R. M. Sibly, The effect of new environment on adapted genetic architecture. *Heredity* **64**, 323–330 (1990).
64. E. G. King, D. A. Roff, D. J. Fairbairn, The evolutionary genetics of acquisition and allocation in the wing dimorphic cricket, *Gryllus firmus*. *Evolution* **65**, 2273–2285 (2011).
65. J. A. Lau, R. G. Shaw, P. B. Reich, P. Tiffin, Indirect effects drive evolutionary responses to global change. *New Phytol.* **201**, 335–343 (2014).
66. F. J. Messina, J. D. Fry, Environment-dependent reversal of a life history trade-off in the seed beetle *Callosobruchus maculatus*. *J. Evol. Biol.* **16**, 501–509 (2003).
67. A. Paccard, M. Vance, Y. Willi, Weak impact of fine-scale landscape heterogeneity on evolutionary potential in *Arabidopsis lyrata*. *J. Evol. Biol.* **26**, 2331–2340 (2013).
68. C. M. Rauter, A. J. Moore, Quantitative genetics of growth and development time in the burying beetle *Nicrophorus pustulatus* in the presence and absence of post-hatching parental care. *Evolution* **56**, 96–110 (2002).
69. R. A. Relyea, The heritability of inducible defenses in tadpoles. *J. Evol. Biol.* **18**, 856–866 (2005).
70. P. Sae-Lim *et al.*, Genotype-by-environment interaction of growth traits in rainbow trout (*Oncorhynchus mykiss*): A continental scale study. *J. Anim. Sci.* **91**, 5572–5581 (2013).
71. P. M. Service, M. R. Rose, Genetic covariation among life-history components: The effect of novel environments. *Evolution* **39**, 943–945 (1985).
72. M. E. Sherrard, H. Maherali, R. G. Latta, Water stress alters the genetic architecture of functional traits associated with drought adaptation in *Avena barbata*. *Evolution* **63**, 702–715 (2009).
73. A. M. Simons, D. A. Roff, The effect of a variable environment on the genetic correlation structure in a field cricket. *Evolution* **50**, 267–275 (1996).
74. A. K. Simonsen, J. R. Stinchcombe, Quantifying evolutionary genetic constraints in the ivyleaf morning glory, *Ipomoea hederacea*. *Int. J. Plant Sci.* **171**, 972–986 (2010).
75. N. Tucic, M. Milosevic, I. Gliksmann, D. Milanovic, I. Aleksic, The effects of larval density on genetic variation and covariation among life-history traits in the bean weevil (*Acanthoscelus obtectus* Say). *Funct. Ecol.* **5**, 525–534 (1991).
76. S. J. Steppan, P. C. Phillips, D. Houle, Comparative quantitative genetics: Evolution of the G matrix. *Trends Ecol. Evol.* **17**, 320–327 (2002).
77. D. Houle, Comparing evolvability and variability. *Genetics* **130**, 195–204 (1992).
78. S. Nakagawa, E. S. A. Santos, Methodological issues and advances in biological meta-analysis. *Evol. Ecol.* **26**, 1253–1274 (2012).
79. W. Viechtbauer, Conducting meta-analyses in R with the metafor package. *J. Stat. Softw.* **36**, 1–48 (2010).
80. D. W. A. Noble, M. Lagisz, R. E. O’dea, S. Nakagawa, Nonindependence and sensitivity analyses in ecological and evolutionary meta-analyses. *Mol. Ecol.* **26**, 2410–2425 (2017).
81. J. P. T. Higgins, S. G. Thompson, Quantifying heterogeneity in a meta-analysis. *Stat. Med.* **21**, 1539–1558 (2002).
82. S. Nakagawa, D. W. A. Noble, A. M. Senior, M. Lagisz, Meta-evaluation of meta-analysis: Ten appraisal questions for biologists. *BMC Biol.* **15**, 18–32 (2017).